

Semiclassical Approach to Finite- N Matrix Models[★]

OLAF LECHTENFELD[†]

*School of Natural Sciences
Institute for Advanced Study
Princeton, NJ 08540*

ABSTRACT

We reformulate the zero-dimensional hermitean one-matrix model as a (non-local) collective field theory, for finite N . The Jacobian arising by changing variables from matrix eigenvalues to their density distribution is treated *exactly*. The semiclassical loop expansion turns out *not* to coincide with the (topological) $\frac{1}{N}$ expansion, because the classical background has a non-trivial N -dependence. We derive a simple integral equation for the classical eigenvalue density, which displays strong non-perturbative behavior around $N = \infty$. This leads to IR singularities in the large- N expansion, but UV divergencies appear as well, despite remarkable cancellations among the Feynman diagrams. We evaluate the free energy at the two-loop level and discuss its regularization. A simple example serves to illustrate the problems and admits explicit comparison with orthogonal polynomial results.

★ Research supported in part by the Department of Energy, contract DE-FG02-90ER40542

† e-mail: OLAF@IASSNS.BITNET or LECHTENF@GUINNESS.IAS.EDU

1. Introduction

Large- N random matrix models have been with us since the 1950s [1], and enjoyed a renaissance 1989 due to newly-found applications to random surfaces, 2d gravity and string theory [2]. In the late 1970s matrix models encountered gauge theories, and its two major calculational frameworks were developed: semiclassical or collective field theory of matrix eigenvalues [3], and the technique of orthogonal polynomials [4]. The latter has driven the recent successes, by admitting the investigation of the double-scaling limit and of the connection to integrable hierarchies. Given that the semiclassical approach is particularly simple in the planar limit, it is somewhat surprising that, except for the one-dimensional model, it has not been extended to the loop level. This paper is intended as an attempt in that direction.

To be sure, there is a well-known obstacle to using the collective field in a $\frac{1}{N}$ expansion of the free energy, namely the exact evaluation of the Jacobian generated by the change of variables from matrix eigenvalues to their density distribution [5-8]. However, recent work of Karabali and Sakita [9] suggests a way of doing just that. In this paper we follow their approach and convert to the Fourier representation of the functional delta-function defining the eigenvalue density $\rho(x)$. Its conjugate field, $\lambda(x)$, gets thereby introduced into an effective action $S_N[\rho, \lambda]$ which, though non-local and non-polynomial, takes on a fairly simple form. An equivalent result has been obtained by Ben-Menahem [10], through mediating the Coulomb repulsion of the eigenvalues by a 2d photon field.

In section 2 we derive the effective action and its equation of motion, for any finite N . The full saddle-point equation contains an explicit $\frac{1}{N}$ term, which obstructs any attempt for an exact solution. At $N = \infty$, it reduces to the airfoil equation, which admits exact solutions vanishing outside a compact support σ [11,12,3]. For $N < \infty$, however, strong non-perturbative (in $\frac{1}{N}$) behavior appears off σ . Section 3 presents the semi-classical loop expansion around the classical background at finite N , *i.e.* one-loop determinant, vertices and propagators. The Feynman graphs for the free energy F_N are drawn at the two-loop level. The exact

but formal expressions are then expanded around $N = \infty$, in section 4. At first we hide the implicit N -dependence of the saddle-point density $\hat{\rho}(x)$ and obtain remarkably simple results for the loop corrections to F_N , due to large cancellations among the Feynman diagrams. We then develop $\hat{\rho}(x)$ in $\frac{1}{N}$ as well by iterating a system of airfoil equations. The $\frac{1}{N}$ expansion is beset by IR and UV singularities whose regularization is briefly discussed. Section 5 works out in detail a simple example and compares to results obtained by orthogonal polynomial techniques. The conclusions form section 6, and some technical material as well as a useful compendium of formulae is collected in three appendices.

2. The Classical Eigenvalue Density

We begin by formulating a collective field theory for the zero-dimensional hermitean one-matrix model at *finite* N . Our starting point is the partition function

$$Z_N \propto \int d^{N^2} M e^{-N \text{tr} V(M)} \quad (2.1)$$

for an $N \times N$ random hermitean matrix ensemble, in a potential V . Upon diagonalization $M = \text{diag}(x_i)$ this reduces to

$$Z_N := \left[\prod_{i=1}^N \int dx_i \right] \exp \left\{ -N \sum_i V(x_i) + \sum_{i < j} \ln(x_i - x_j)^2 \right\} \quad (2.2)$$

We like to change variables from the matrix eigenvalues x_i to their density distribution[★]

$$\rho(x) := \frac{1}{N} \sum_{i=1}^N \delta(x - x_i) \quad (2.3)$$

★ Obviously, this transformation is $N : \aleph_1$ but then ρ vanishes almost everywhere.

More precisely, we insert

$$\begin{aligned}
1 &= \int \mathcal{D}\rho \prod_x \delta\left(\rho(x) - \frac{1}{N} \sum_i \delta(x-x_i)\right) \\
&= \iint \mathcal{D}\rho \mathcal{D}\lambda \exp\left\{i \int dx \lambda(x) \left[\rho(x) - \frac{1}{N} \sum_i \delta(x-x_i)\right]\right\}
\end{aligned} \tag{2.4}$$

into (2.2) and express the action in terms of the density,

$$S_N^0[\rho] = N^2 \int dx \rho(x) V(x) - \frac{1}{2} N^2 \iint dx dy \rho(x) f(x-y) \rho(y) + \frac{1}{2} N f(0) \quad . \tag{2.5}$$

The self-interaction had to be regulated by replacing $\ln z^2 \rightarrow f(z)$ in eq. (2.2), choosing some suitable, *i.e.* symmetric and bounded, function f .

Following ref. 9 we are able to perform the integration over x_i ,

$$\begin{aligned}
Z_N &= \iint \mathcal{D}\rho \mathcal{D}\lambda \ e^{-S_N^0[\rho] + i \int \lambda \rho} \left[\prod_{i=1}^N \int dx_i \right] e^{-\frac{i}{N} \sum_i \lambda(x_i)} \\
&= \iint \mathcal{D}\rho \mathcal{D}\lambda \ e^{-S_N^0[\rho] + i \int \lambda \rho} \left[\int dx \ e^{-\frac{i}{N} \lambda(x)} \right]^N ,
\end{aligned} \tag{2.6}$$

and arrive at an effective action

$$S_N[\rho, \lambda] = S_N^0[\rho] - i \int dx \lambda(x) \rho(x) - N \ln \int dx e^{-\frac{i}{N} \lambda(x)} \tag{2.7}$$

which is not only nonlocal in the two real fields ρ and λ but also non-polynomial in the latter. Interestingly, the constant mode of λ can be integrated out exactly to yield the constraint $\delta(\int \rho - 1)$ that was apparent already from the definition (2.3). However, we shall keep those modes in the measure for the time being. In principle, another constraint arises from the positivity of ρ . Perturbation theory about a strictly positive $\hat{\rho}$, however, is insensitive to this restriction, and we will therefore ignore it in the fluctuations.

Our goal is to carry out a systematic semiclassical analysis of this peculiar one-dimensional field theory. To leading order in $\hbar$ we must determine the saddle-point configurations $(\hat{\rho}, \hat{\lambda})$, where the action (2.7) is stationary. The first variations yield

$$\begin{aligned} 0 &= N^2 V(x) - N^2 \int dy f(x-y) \hat{\rho}(y) - i \hat{\lambda}(x) \\ 0 &= \hat{\rho}(x) - e^{-\frac{i}{N} \hat{\lambda}(x)} \bigg/ \int dy e^{-\frac{i}{N} \hat{\lambda}(y)} \end{aligned} \quad (2.8)$$

where $\hat{\rho}$ comes out to be properly normalized, $\int \hat{\rho} = 1$. The second equation determines $\hat{\lambda}$ up to a constant,

$$\hat{\lambda}(x) = iN \ln \hat{\rho}(x) + \lambda_0 \quad , \quad (2.9)$$

a result which may be inserted into the first equation.[★] Subtracting the value at $x = 0$ to get rid of the constant and deregulating the Coulomb repulsion we get[†]

$$[V(x) - V(0)] - \int dy [\ln(x-y)^2 - \ln y^2] \hat{\rho}(y) + \frac{1}{N} [\ln \hat{\rho}(x) - \ln \hat{\rho}(0)] = 0 \quad . \quad (2.10)$$

Together with normalization and positivity, this equation describes the classical eigenvalue density for any finite N . It is noteworthy that (2.10) is not homogeneous in $\frac{1}{N}$, so its solution cannot be, either. The saddle-point equation is more easily studied in its differentiated version,

$$\frac{1}{2} V'(x) + \frac{1}{2N} \frac{\hat{\rho}'}{\hat{\rho}}(x) = \oint \frac{dy}{x-y} \hat{\rho}(y) \equiv \pi \mathcal{H}_x[\hat{\rho}] \quad (2.11)$$

where $\oint$ denotes Cauchy's principal value of the integral. The *r.h.s.* is known as the Hilbert transform $\mathcal{H}$ (of $\hat{\rho}$) which has been thoroughly investigated [13] (see

★ The first equation subsequently fixes the constant λ_0 for a given solution $\hat{\rho}$.

† See also ref. [10].

appendix A for details). At this point let us merely state that $\mathcal{H}$ is uniquely invertible on the class L^p , $p > 1$, by

$$\mathcal{H}_x^{-1}[\phi] = -\frac{1}{\pi} \oint \frac{dy}{x-y} \phi(y) \quad . \quad (2.12)$$

At $N = \infty$ our equation (2.11) has been widely studied and solved [11,12,3], and it was learned [14] that a *unique* solution extends to $N < \infty$. Unfortunately, the equation is not easily solved for finite N . Even in the large- N limit some care is required, as shown by the following. For instance, should we drop the $\hat{\rho}'/\hat{\rho}$ -term since it is down by $\frac{1}{N}$? A little inspection reveals that such a step is in general not consistent with the asymptotic large- $|x|$ behavior of the equation, which demands a $\frac{1}{x}$ fall-off for the *l.h.s.* due to the normalization of $\hat{\rho}$. In fact, the formal iterations

$$\begin{aligned} \hat{\rho}(x) &= -\frac{1}{\pi^2} \oint \frac{dy}{x-y} \left[\frac{1}{2} V'(y) + \frac{1}{2N} \frac{\hat{\rho}'}{\hat{\rho}}(y) \right] \\ \hat{\rho}(x) &\sim \exp \left\{ -N \left[V(x) - \int \ln(x-y)^2 \hat{\rho}(y) \right] \right\} \stackrel{|x| \rightarrow \infty}{\approx} x^{2N} e^{-NV(x)} \end{aligned} \quad (2.13)$$

show all terms in (2.11) to be of the same order for $|x| \gg 1$ (unless $V \sim \ln x^2$). The situation is different, however, near the minimum of the potential where most of $\hat{\rho}$ is concentrated.[‡] When $\hat{\rho}$ is $\mathcal{O}(1)$ the $\hat{\rho}'/\hat{\rho}$ -term may safely be neglected or treated as a $\frac{1}{N}$ perturbation in (2.11). To the same approximation one should also drop the exponential tail of $\hat{\rho}$ under the integral. At $N = \infty$ this amounts to setting $\hat{\rho}$ identically to zero beyond a certain distance from the minimum of V . The cross-over points $x = a, b$ between the exterior region (where $\hat{\rho}$ is $\mathcal{O}(e^{-N})$) and the interior domain $\sigma \equiv (a, b)$ (where it is $\mathcal{O}(1)$) are determined by the normalization of $\hat{\rho}$; they occur where $\mathcal{H}[\hat{\rho}]$ can no longer match the growth of V' . For $N \rightarrow \infty$, $\hat{\rho} \rightarrow \rho_0$, it is therefore indeed consistent to simplify eq. (2.11) to [3]

$$\begin{aligned} \frac{1}{2} V'(x) &= \oint_a^b \frac{dy}{x-y} \rho_0(y) \equiv \pi \mathcal{T}_x[\rho_0] & \text{for } x \in (a, b) \\ \rho_0(x) &= 0 & \text{for } x \notin (a, b) \quad . \end{aligned} \quad (2.14)$$

[‡] For simplicity we restrict ourselves to single-well potentials. See, however, ref. 14.

This relation is known as the *airfoil equation* and determines the vorticity ρ_0 related to a given velocity field V' along the airfoil σ ($b-a$ is the span of the wings) [11]. It is solved by inverting the *finite* Hilbert transform $\mathcal{T}$,

$$\rho_0(x) = \frac{1}{\sqrt{(b-x)(x-a)}} \left[\gamma - \frac{1}{\pi^2} \int_a^b \frac{dy}{x-y} \frac{1}{2} V'(y) \sqrt{(b-y)(y-a)} \right] \quad (2.15)$$

where the arbitrary constant

$$\gamma = \frac{1}{\pi} \int_a^b dy \rho_0(y) \quad (2.16)$$

arises due to the one-dimensional kernel $\frac{1}{\sqrt{(b-x)(x-a)}}$ of $\mathcal{T}$ and is set to $\gamma = \frac{1}{\pi}$ by the normalization of ρ_0 . The inversion formula (2.15) is guaranteed to produce a solution to (2.14) as long as $V' \in L^{\frac{4}{3}+\epsilon}$; for $V' \in L^{1+\epsilon}$ we also need $\rho_0 \in L^{1+\epsilon}$ in (2.15) [11, 12]. In the case of polynomial potentials the resulting density $\rho_0(x)$ is seen to vanish at the endpoints of σ generically as $(x-a)^{1/2}$ and $(b-x)^{1/2}$. For more on the finite Hilbert transform we refer to appendix A.

Having solved the saddle-point equation in the large- N limit, one is tempted to try and expand the complete solution $\hat{\rho}$ of (2.11) around ρ_0 . A $\frac{1}{N}$ -expansion appears logical on σ but is bound to fail in the exterior region since e^{-N} has an essential singularity at $N = \infty$. Instead, a series in powers of e^{-NV} seems more appropriate there. Either type of expansion, however, must break down at the cross-over points, because a smooth function $\hat{\rho}$ is to develop from one that is singular there. For finite N the transition from $\mathcal{O}(1)$ to $\mathcal{O}(e^{-N})$ occurs in a finite intermediate region of size $N^{-2/3}$ by a generalized Airy function of the scaling variable $(x-b)N^{2/3}$. Still, one might hope that a $\frac{1}{N}$ -expansion of the classical action needs $\hat{\rho}$ only on σ , due to its non-perturbative nature off σ . We shall pursue this line of argument in Sect. 4.

It remains to evaluate the classical action in terms of $\hat{\rho}$. Employing eq. (2.9) as well as $\ln \int e^{-\frac{i}{N}\hat{\lambda}} = -\frac{i}{N}\lambda_0$ we find, in agreement with ref. [10], that

$$S_N[\hat{\rho}, \hat{\lambda}] = N^2 \int \hat{\rho} V - \frac{1}{2} N^2 \iint \hat{\rho} f \hat{\rho} + N \int \hat{\rho} \ln \hat{\rho} + \frac{1}{2} N f(0) \quad (2.17)$$

which may be rewritten as

$$\begin{aligned} S_N[\hat{\rho}, \hat{\lambda}] &= \frac{N^2}{2} \int \hat{\rho} (V-f) + \frac{N^2}{2} V(0) + \frac{N}{2} \int \hat{\rho} \ln \hat{\rho} + \frac{N}{2} \ln \hat{\rho}(0) + \frac{N}{2} f(0) \\ &= \frac{N^2}{2} \iint \hat{\rho}(x) [f(x-y) - f(x) - f(y)] \hat{\rho}(y) + N \ln \hat{\rho}(0) + \frac{N}{2} f(0) \end{aligned} \quad (2.18)$$

by making use of the saddle-point equation (2.10). Given the non-trivial N -dependence of $\hat{\rho}$ it becomes clear that the semiclassical expansion will not correspond exactly to an expansion in $\frac{1}{N}$. On the other hand, it is encouraging to note that potentially divergent pieces like $\int \ln \hat{\rho}$ have dropped out. The eventually singular $f(0)$ term will get cancelled in the following section.

3. Quantum Fluctuations

We now set out to develop the loop expansion of the theory defined by eqs. (2.5) and (2.7) in the standard fashion. We eliminate the constant modes from the measure and split

$$\rho(x) = \hat{\rho}(x) + \rho'(x) \quad \lambda(x) = \hat{\lambda}(x) + \lambda'(x) \quad (3.1)$$

with $\int \rho' = 0 = \int \lambda'$. Since S_N is quadratic in ρ its fluctuations are gaussian and affect only the propagators. Hence, only vertices for λ' are present. Let us write

$$Z_N = Z_{\text{cl}} Z_{1\text{-loop}} Z_+ \quad Z_{\text{cl}} = e^{-S_N[\hat{\rho}, \hat{\lambda}]} \quad Z_{1\text{-loop}} = [\det' S_N'']^{-\frac{1}{2}} \quad (3.2)$$

where $\ln Z_+$ equals the connected Feynman diagrams, and $\det'$ reminds us to omit

the constant modes. The fluctuation operator S''_N at the saddle point reads

$$S''_N(x, y) = \begin{pmatrix} -N^2 f(x-y) & -i\delta(x-y) \\ -i\delta(x-y) & \frac{1}{N} [\hat{\rho}(x) \delta(x-y) - \hat{\rho}(x) \hat{\rho}(y)] \end{pmatrix} \quad (3.3)$$

and acts on $\{\rho'(y), \lambda'(y)\}$. The simple form of the off-diagonal blocks of S''_N facilitates the calculation of the determinant,[★]

$$\begin{aligned} Z_{1\text{-loop}} &= \det' \left[\delta(x-y) - N f(x-y) \hat{\rho}(y) + N \int dz f(x-z) \hat{\rho}(z) \hat{\rho}(y) \right]^{-\frac{1}{2}} \\ &= \exp \left\{ \sum_{k=1}^{\infty} \frac{N^k}{2k} \text{tr} \left[f(x-y) \hat{\rho}(y) - \int dz f(x-z) \hat{\rho}(z) \hat{\rho}(y) \right]^k \right\}, \end{aligned} \quad (3.4)$$

where we have formally expanded in powers of N (not $\frac{1}{N}$). In the obvious graphical representation this gives rise not only to the usual loop diagrams but also to loops “cut open” at any line, due to the second term in the trace. Each term in the above expansion seems to be well-behaved except for the very first one, $\frac{N}{2} \text{tr} f \hat{\rho} = \frac{N}{2} f(0)$, which in fact just cancels the eventually singular term in the classical action, eq. (2.17). Of course, one would rather like to expand in powers of $\frac{1}{N}$ but let us defer this to the next section.

Instead, we complete the construction of the Feynman rules. To obtain the λ' propagator we need to invert S''_N ,[†]

$$\begin{aligned} \lambda'(x) \lambda'(y) &= N \left[(\hat{\rho}(x) \delta(x-y) - \hat{\rho}(x) \hat{\rho}(y)) - \frac{1}{N} f^{-1}(x, y) \right]^{-1} \\ &= -N^2 f(x, y) - N^3 \iint f(x, z) (\hat{\rho}(z) \delta(z-w) - \hat{\rho}(z) \hat{\rho}(w)) f(w, y) - \dots \end{aligned} \quad (3.5)$$

again formally expanding in N . The corresponding graphs are the same as in (3.4) before taking the trace.

★ $\det \begin{pmatrix} A & -i\mathbf{1} \\ -i\mathbf{1} & B \end{pmatrix} = \det [\mathbf{1} + A \cdot B]$. Since 1 is a zero mode of B we may omit the primes.

† The inversion of f will be discussed in the next section.

Turning to the vertices, the λ' expansion simplifies at the saddle point as

$$\begin{aligned}\Delta S &\equiv -N \ln \int dx e^{-\frac{i}{N}\lambda(x)} = i\lambda_0 - N \ln \int dx \hat{\rho}(x) e^{-\frac{i}{N}\lambda'(x)} \\ &= i\lambda_0 + N \sum_{r=1}^{\infty} \frac{(-1)^r}{r} \left[\sum_{s=1}^{\infty} \frac{1}{s!} \left(\frac{-i}{N} \right)^s \int dx \hat{\rho}(x) \lambda'(x)^s \right]^r\end{aligned}\quad (3.6)$$

with the help of $e^{-\frac{i}{N}\hat{\lambda}} / \int e^{-\frac{i}{N}\hat{\lambda}} = \hat{\rho}$. The constant mode of λ' appears only at $\mathcal{O}(1)$, as it should. Note that this is a genuine $\frac{1}{N}$ expansion because in powers of $\frac{i}{N}\lambda'$. The $\mathcal{O}(1)$ and $\mathcal{O}(\frac{1}{N})$ terms have already been taken care of in (2.8) and (3.3).

It is instructive to compute the two-loop correction to the free energy

$$F_N = -\ln Z_N = S_N[\hat{\rho}, \hat{\lambda}] + \frac{1}{2} \text{tr}' \ln S_N'' - \ln Z_+ \quad . \quad (3.7)$$

Writing $\Delta S = N \sum_k N^{-k} S_k$, with k lines emanating from S_k , we have

$$Z_+ = \left\langle e^{-S_3/N^2 - S_4/N^3 - \dots} \right\rangle = \exp \left\{ -N^{-3} \langle S_4 \rangle + \frac{1}{2} N^{-4} \langle S_3 S_3 \rangle - \dots \right\} \quad (3.8)$$

where $\langle \dots \rangle$ denotes the free field average performed with the propagator (3.5).

With a little combinatorics S_k can be read off eq. (3.6), *e.g.*

$$\begin{aligned}S_3 &= \frac{(-i)^3}{3!} \left[-\int \hat{\rho} \lambda'^3 + 3 \int \hat{\rho} \lambda'^2 \int \hat{\rho} \lambda' - 2 (\int \hat{\rho} \lambda')^3 \right] \\ S_4 &= \frac{(-i)^4}{4!} \left[-\int \hat{\rho} \lambda'^4 + 4 \int \hat{\rho} \lambda'^3 \int \hat{\rho} \lambda' + 3 (\int \hat{\rho} \lambda'^2)^2 - 12 \int \hat{\rho} \lambda'^2 (\int \hat{\rho} \lambda')^2 + 6 (\int \hat{\rho} \lambda')^4 \right]\end{aligned}\quad (3.9)$$

which are depicted in figure 1. Obviously, the parts of S_k are labelled by the partitions of the number k into integers. The λ' contractions produce 21 diagrams which are displayed in figure 2. Due to the non-locality of the vertices, the terms “loop” and “vertex” take on *two* distinct meanings, namely one counting powers of $\hbar$ and denoting S_k , and the other describing individual diagrams. In order to

avoid notational confusion, we distinguish between the two and refer to figure 2 as the two-loop *contributions* built from *vertices* S_k , represented by 6 two-loop, 9 one-loop, and 6 tree-level *diagrams* consisting of 2 or 3 lines and up to 6 *nodes*. Note also that all contributions to $\ln Z_+$ are connected whereas the constituting graphs may not be. It appears that we have derived a sensible set of Feynman rules (3.4)-(3.6) for our non-local theory.

The functional ΔS in (3.6) has the property that an x -independent shift of λ' leaves the vertices S_k form-invariant. We may exploit this fact to simplify the structure of S_k , by changing the field variable $\lambda' \rightarrow \tilde{\lambda} := \lambda' - \int \hat{\rho} \lambda'^*$. Since $\int \hat{\rho} \tilde{\lambda} = 0$, a large number of terms disappear when S_k are expressed in terms of $\tilde{\lambda}$. We are left with the parts corresponding to partitions of k into integers ≥ 2 only. The 21 diagrams of figure 2 compactify to the five graphs without univalent nodes. In particular, tree graphs can no longer be present. The price we pay for this simplification is a more involved propagator for $\tilde{\lambda}$, which actually consists of four terms when rewritten in terms of the λ' propagator.

We gain, however, an argument showing the cancellation of all tadpole infinities. The only potentially divergent diagrams arise from self-contractions,

$$\begin{array}{c} \lambda'(x) \lambda'(x) \\ \boxed{} \end{array} = -N^2 f(0) + \text{finite terms} \quad . \quad (3.10)$$

Any $\tilde{\lambda}$ -propagator ending in a univalent node with a weight of $\text{constant} \times \hat{\rho}$ is seen to vanish, which eliminates all $f(0)$ terms from tadpoles. The singularities arising from self-contracted multiply valent nodes appear to cancel systematically between different graphs (see figure 2 for example). This argument holds true for any *constant* part of $\lambda'(x) \lambda'(x)$, whether we expand in N or not.

Expanding the full propagators in the loop graphs à la (3.5) leads to an infinite series in powers of N , which ruins the $\frac{1}{N}$ expansion from (3.6). Indeed,

★ The Jacobian of this transformation is compensated by the change in the one-loop determinant.

absorbing the leading N -dependence of $\lambda'(x)\lambda'(y)$ into the vertices in (3.8) replaces $S_k/N^{k-1} \rightarrow NS_k$, for any k , so that all terms in ΔS contribute at lowest order in N . It therefore appears that we need to sum the loop expansion in order to consistently develop F_N in powers of N . On the other hand, F_N has a logarithmic singularity at $N=0$ (at least for a quadratic matrix potential) which makes an N expansion ill-defined anyway.

Nevertheless, the semiclassical expansion seems to be well-behaved for finite N . It is merely dangerous to develop the individual loop contributions in N (or $\frac{1}{N}$ as we shall see soon). Unfortunately, for practical purposes we need to compute the determinant (3.4) and the propagator (3.5) somewhere, and this seems feasible only at the extrema $N=0$ or $N=\infty$. Another option is to put the system in a box (in eigenvalue space) of length L and estimate determinant and propagator in Fourier space. We shall come back to this approach later on.

4. $\frac{1}{N}$ Expansion

This section is devoted to an attempt to set up a systematic $\frac{1}{N}$ expansion of the free energy (3.7), using the previous results (2.11)-(2.18) and (3.4)-(3.6). We will proceed in two stages. First, we arrange according to explicit powers of $\frac{1}{N}$ and hide the implicit N -dependence of $\hat{\rho}$. Second, we expand $\hat{\rho}$ as well and insert the result into the coefficients obtained earlier.

The classical part of the free energy given by eqs. (2.17) or (2.18), it remains to develop the fluctuation determinant (3.4) and the full λ' propagator (3.5) around $N=\infty$. To this end one needs to invert $NS_N''|_{\lambda\lambda} = \hat{\rho}(x)\delta(x-y) - \hat{\rho}(x)\hat{\rho}(y)$ away from the constant mode. As expected, this operator has a constant zero mode since $\int \hat{\rho} = 1$. Hence, its inverse orthogonal to that mode is simply[†]

$$[\hat{\rho}(x)\delta(x-y) - \hat{\rho}(x)\hat{\rho}(y)]_{\perp}^{-1} = \frac{1}{\hat{\rho}(x)} \delta(x-y) \quad . \quad (4.1)$$

[†] It is convenient to make a change of basis, $\lambda' = \xi/\sqrt{\hat{\rho}}$, so that the kernel of $\delta(x-y) - \sqrt{\hat{\rho}(x)\hat{\rho}(y)}$ is spanned by $\sqrt{\hat{\rho}}$. Because $\sqrt{\hat{\rho}(x)\hat{\rho}(y)}$ projects onto the kernel, our operator projects onto the complement and is inverted by the identity there.

On the level of the determinant this can also be seen in Fourier space, from the identity

$$\det'[\hat{\rho}_{k+l} - \hat{\rho}_k \hat{\rho}_l] = \det[\hat{\rho}_{k+l}] \quad \text{for} \quad \hat{\rho}_0 = 1 \quad . \quad (4.2)$$

The propagator (3.5) is then easily written out as

$$\underbrace{\lambda'(x) \lambda'(y)} = N \frac{1}{\hat{\rho}(x)} \delta(x-y) + \frac{1}{\hat{\rho}(x)} f^{-1}(x, y) \frac{1}{\hat{\rho}(y)} + \frac{1}{N} \int \frac{1}{\hat{\rho}} f^{-1} \frac{1}{\hat{\rho}} f^{-1} \frac{1}{\hat{\rho}} + \mathcal{O}(\frac{1}{N^2}) . \quad (4.3)$$

The one-loop determinant (3.4) works out analogously,

$$\begin{aligned} Z_{1\text{-loop}} &= \det \left[-N f(x-y) \hat{\rho}(y) \right]^{-\frac{1}{2}} \times \det \left[\delta(x-y) - \frac{1}{N} \frac{1}{\hat{\rho}(x)} f^{-1}(x, y) \right]^{-\frac{1}{2}} \\ &\propto \exp \left\{ -\frac{1}{2} \eta \ln N - \frac{1}{2} \int \ln \hat{\rho} \right\} \times \exp \left\{ \sum_{k=1}^{\infty} \frac{N^{-k}}{2k} \text{tr} \left[\hat{\rho}(x)^{-1} f^{-1}(x, y) \right]^k \right\} \end{aligned} \quad (4.4)$$

where η is a regulating constant, and we dropped a singular constant determinant. Together with eqs. (2.18) and (3.6) this yields a systematic but formal expansion of the free energy (3.7). Each k -loop contribution starts at $\mathcal{O}(N^{-k+1})$ and is made of k - and lesser-loop graphs, many of which are disconnected. The ensuing Feynman rules are given in figure 3, but the combinatorial coefficients and the occurrence of the nodes have to be read off the expansions (3.6), (4.3) and (4.4). Clearly, the expansion is not just in $\frac{1}{N}$ but also in $\frac{1}{N\hat{\rho}}$, which will be responsible for IR divergencies later on.

We have yet to give a less formal meaning to the “propagator” f^{-1} . Since the convolution with $\frac{1}{2\pi} \partial_x f(x, y) \rightarrow \frac{1}{\pi} \text{P} \frac{1}{x-y}$ is just the Hilbert transform (2.11), the deregulated $f(x-y) \rightarrow \ln(x-y)^2$ is easily inverted from (2.12),[‡]

$$f^{-1}(x, y) \rightarrow -\frac{1}{2\pi^2} \text{P} \frac{1}{x-y} \frac{d}{dy} = -\frac{1}{2\pi^2} \frac{d}{dx} \text{P} \frac{1}{x-y} \quad (4.5)$$

on the space $L^{1+\epsilon}$.

[‡] The symbol P denotes the principal value, $\int \text{P} \dots = \oint \dots$, or $\text{P} \frac{1}{x} = \frac{1}{x} - i\pi \delta(x)$.

Employing the results of appendix A we are now able to compute the one-loop traces of eq. (4.4). The derivatives get distributed over the graph, and, after evaluating

$$\begin{aligned} \iint \frac{P}{y-x} \phi_1(x) \frac{P}{x-y} \phi_2(y) &= \int \mathcal{H}'[\phi_1] \phi_2 - \delta(0) \int \phi_1 \phi_2 \\ \iiint \frac{P}{z-x} \phi_1(x) \frac{P}{x-y} \phi_2(y) \frac{P}{y-z} \phi_3(z) &= \int (\mathcal{H}'[\phi_1] \phi_2 - \phi_1 \mathcal{H}'[\phi_2]) \mathcal{H}[\phi_3] \end{aligned} \quad (4.6)$$

and so on, we discover that $\text{tr}[\hat{\rho}^{-1} f^{-1}]^k$ vanish upon symmetrization over cyclic permutations![§] Hence, the one-loop correction to F_N comes from the first (singular) factor of (4.4) only and is given by

$$\frac{1}{2} \text{tr}' \ln S_N'' = \frac{1}{2} \eta \ln N + \frac{1}{2} \int dx \ln \hat{\rho}(x) \quad . \quad (4.7)$$

By the same calculation, self-contractions $\underbrace{\lambda'(x) \lambda'(x)}_{\text{self-contraction}}$ do not receive $\frac{1}{N}$ corrections beyond the leading term.

To order N^{-1} we have to review our two-loop calculation of the previous section. It suffices to replace all full propagators of figure 2 by $N \hat{\rho}(x)^{-1} \delta(x-y)$. In terms of the Feynman rules of figure 3 this means that we compose our diagrams with dashed lines only. Thus any propagator loop yields a singular factor of $\delta(0)$, leaving us with connected graph components proportional to $\delta(0)^2 \hat{\rho}^{-1}$, $\delta(0)$, or $\hat{\rho}$. It is quite remarkable that all but one of the divergent graphs cancel between $\langle S_4 \rangle$ and $\langle S_3 S_3 \rangle$![¶] However, the single genuine two-loop diagram survives (together with some finite contributions),

$$-\ln Z_+ = \left[\frac{1}{12} \delta(0)^2 \int \frac{dx}{\hat{\rho}(x)} - \frac{1}{12} \right] \frac{1}{N} + \mathcal{O}\left(\frac{1}{N^2}\right) \quad . \quad (4.8)$$

It is then quite clear that a k -loop contribution to $\ln Z_+$ will yield (products of) expressions of the form $\delta(0)^p \int \hat{\rho}^{1-p}$, with $0 \leq p \leq k$. However, all Feynman graphs

[§] We have checked this for $k \leq 3$.

[¶] The propagator self-contraction cancellation argument of the previous section does not even apply since $\underbrace{\lambda'(x) \lambda'(x)}_{\text{self-contraction}} = N \delta(0) / \hat{\rho}(x)$ is not constant.

appear to cancel each other except for those with $p=0$ and $p=k$. The latter are simply given by all genuine (connected) k -loop graphs, presumably without self-contractions.*

It should not have escaped the attentive reader that the expressions (4.7) and (4.8) are divergent. In fact, we encounter two different types of singularities. First, the regulator η as well as the factors of $\delta(0)$ arose from evaluating $\text{tr } \mathbf{1}$ and thus count the (infinite) number of degrees of freedom. Second, the integrals in both expressions are singular since $\hat{\rho} \sim e^{-NV}$ for $|x| \rightarrow \infty$ induces $\int \ln \hat{\rho} \sim N \int V$ and $\int \hat{\rho}^{-1} \sim \int e^{+NV}$. The latter infinities require an IR cutoff; most convenient is a restriction of the eigenvalue range to a finite interval, say $x \in [-\frac{L}{2}, +\frac{L}{2}]$, with open boundary conditions. Putting the system into a box does not, however, cure the UV singularities mentioned first; those ask for a mode cutoff, say $|k| \leq M$, for some large M . A natural choice for the value of M would seem $M = \alpha N + \beta$. We have estimated the resulting leading behavior of the one-loop determinant (3.4), with some details given in appendix C. The upshot is

$$\frac{1}{2} \text{tr}' \ln S_N'' \Big|_{\text{reg}} = [(\alpha+1) \ln(\alpha+1) - \alpha \ln \alpha] N + \mathcal{O}(\ln N) \quad . \quad (4.9)$$

The $\mathcal{O}(N)$ term originates from $\frac{1}{2} \ln \det[-f]$ which we dropped in (4.4). Since we did not probe the $\hat{\rho}$ dependence, L does not appear in our result. Further numerical analysis is required to elucidate the cut-off dependence of the regulated loop corrections.

The coefficients of the $\frac{1}{N}$ series we have obtained are (singular) functionals of the saddle-point density $\hat{\rho}$. To complete the $\frac{1}{N}$ expansion we finally develop $\hat{\rho}$ about $N = \infty$ as well, thus blinding ourselves to its “non-perturbative” parts. In $\frac{1}{N}$ perturbation we set $\hat{\rho}(x) \equiv 0$ for $x \notin \sigma$ and expand

$$\hat{\rho}(x) = \sum_{i=0}^{\infty} N^{-i} \rho_i(x) \quad \text{for } x \in \sigma \quad . \quad (4.10)$$

* We have checked this also on the 3-loop level, with an outcome of $\frac{1}{12} \delta(0)^3 \int (\hat{\rho} N)^{-2} + \text{finite}$.

Next, we define a sequence of potentials

$$V_0 = V \quad , \quad V_1 = \ln \rho_0 \quad , \quad \sum_{i=1}^{\infty} N^{-i} V_{i+1} = \sum_{r=1}^{\infty} \frac{(-)^{r-1}}{r} \left(\sum_{s=1}^{\infty} N^{-s} \frac{\rho_s}{\rho_0} \right)^r \quad (4.11)$$

by expanding $\ln \hat{\rho}$. The saddle-point equation (2.11) then turns into the iteration

$$\frac{1}{2} V'_i(x) = \int_a^b \frac{dy}{x-y} \rho_i(y) \quad \text{for } x \in (a, b) \quad (4.12)$$

since V_i is a function of $\{\rho_0, \rho_1, \dots, \rho_{i-1}\}$. The constant part of ρ_i is fixed by the normalization conditions $\int \rho_i = \delta_{i,0}$. Clearly, the iteration consists of repeated use of eqn. (2.15) for $\mathcal{T}^{-1}$, with $\gamma=0$ after the first step.

Having computed the coefficients in (4.10) to a desired order in $\frac{1}{N}$, we are able to unravel the complete large- N expansion of the free energy. First, the above equations can be employed to simplify the $\frac{1}{N}$ series of the classical action (2.18):

$$\begin{aligned} S_N[\hat{\rho}, \hat{\lambda}] &= \frac{N^2}{2} \int \rho_0 [V(x) - V(0) - f(x)] + \frac{N}{2} f(0) + N \int \rho_0 \ln \rho_0 + \frac{1}{2} \int \rho_1 \ln \rho_0 + \dots \\ &= \frac{1}{2} N^2 \iint \rho_0(x) [f(x-y) - f(x) - f(y)] \rho_0(y) + \frac{1}{2} N f(0) + N \ln \rho_0(0) \\ &\quad + N \iint \rho_0(x) [f(x-y) - f(y)] \rho_1(y) + \frac{1}{2} \iint \rho_1(x) f(x-y) \rho_1(y) + \mathcal{O}(\frac{1}{N}) . \end{aligned} \quad (4.13)$$

Second, we insert the expansion (4.10) into the expressions (4.7) and (4.8) for the loop corrections. However, by setting $\hat{\rho} \sim e^{-NV}$ to zero off σ we are dropping infinite contributions $\sim N \int V$ and $\sim \int e^{+NV}$ to the integrals of (4.7) and (4.8), respectively, of which the first is perfectly perturbative.

At this stage it is quite obvious that a brute-force expansion around $N = \infty$ makes no sense. In view of the interference with strong e^{-N} behavior one must keep at least the N -dependence of $\hat{\rho}$ intact. Only then may one hope to establish a large- N series for the free energy, provided the regulation problems can

be overcome. Under these circumstances it is hard to see how a serious comparison with orthogonal polynomial results could be achieved beyond the leading order.

5. An Example

We have tried to develop the semiclassical expansion for the zero-dimensional hermitean matrix model with an arbitrary potential V . In order to explicitly demonstrate the difficulties of this scheme we now become more specific and choose a simple potential,

$$V(x) = \frac{1}{2}x^2 + g x^4 \quad , \quad g > 0 \quad , \quad (5.1)$$

for which we can evaluate the free energy F_N by the method of orthogonal polynomials. From the literature [4] we extract the value for $F_N(g) - F_N(0)$, and add to it the result for the quadratic potential ($g=0$), as calculated in appendix B. The first orders in $\frac{1}{N}$ yield

$$\begin{aligned} F_N = & \left[\frac{3}{4} + \frac{1}{24}(a^2 - 1)(9 - a^2) - \frac{1}{2} \ln a^2 \right] N^2 - N \ln N \\ & + \left[1 - \ln 2\pi \right] N - \frac{5}{12} \ln N + \left[c_0 + \frac{1}{12} \ln(2 - a^2) \right] + \mathcal{O}(N^{-2}) \end{aligned} \quad (5.2)$$

with

$$12ga^4 + a^2 - 1 = 0 \quad \Longleftrightarrow \quad a^2 = (-1 + \sqrt{1 + 48g})/24g \quad (5.3)$$

and $c_0 \approx -0.7535$, a universal constant.

Let us see how much of this the collective field is able to reproduce. The iterative solution of eqs. (4.11) and (4.12) proceeds along standard lines. To lowest order eq. (2.15) produces [3]

$$\rho_0(x) = \frac{1}{2\pi} (1 + 8ga^2 + 4gx^2) \sqrt{4a^2 - x^2} \Theta(4a^2 - x^2) \quad , \quad (5.4)$$

a polynomial deformation of Wigner's semicircle law. The next step in the itera-

tion (4.11) and (4.12) yields

$$V_1'(x) = \frac{2x}{x^2 + n^2} - \frac{x}{4a^2 - x^2} \quad \text{with} \quad n^2 = \frac{1+8ga^2}{4g} = \frac{2+a^2}{1-a^2}a^2 \quad (5.5)$$

which is to be plugged into (2.15) to find the correction $\rho_1 = \rho_1^{(1)} + \rho_1^{(0)}$. With the help of appendix A^{*} the first term gives

$$\rho_1^{(1)}(x) = \frac{1}{\pi} \frac{1}{\sqrt{4a^2 - x^2}} \left[\frac{\sqrt{(4a^2 + n^2)n^2}}{x^2 + n^2} - 1 \right] \Theta(4a^2 - x^2) \quad (5.6)$$

but the second term is not integrable at $x = \pm 2a$; stretching the meaning of (4.12) we might set

$$\rho_1^{(0)}(x) = \frac{1}{4} [\delta(x - 2a) + \delta(x + 2a)] - \frac{1}{4\pi} \frac{\Theta(4a^2 - x^2)}{\sqrt{4a^2 - x^2}}. \quad (5.7)$$

These results continue to hold true for $-\frac{1}{48} < g < 0$; in this regime $2 > a^2 > 1$ and $-8 > n^2 > -\infty$. At the lower bound $|n| \rightarrow 2a$, and the well-known additional zero develops at the edge of σ . Evidently, the iteration (4.12) leads to ever worse singularities at $x = \pm 2a$. This merely reflects the breakdown of the $\frac{1}{N}$ expansion at the edges and shows that our neglect of the non-perturbative tail of $\hat{\rho}$ was illegitimate.

Nevertheless, we insert the above findings into the functionals (4.13), (4.7) and (4.8). The classical action becomes

$$\begin{aligned} S_N[\hat{\rho}, \hat{\lambda}] &= N^2 \left[\frac{3}{4} + \frac{1}{24}(a^2 - 1)(9 - a^2) - \frac{1}{2} \ln a^2 \right] + \frac{1}{2} N f(0) \\ &+ N \left[-\ln \pi + a^2 \left(\frac{1}{2} - \ln 2 \right) + \ln \frac{2+a^2}{3a} + (1-a^2) \left(\frac{5}{12} - \ln 2 \right) \right. \\ &\quad \left. + 2 \ln \frac{1}{2}(1+b) + \frac{1}{3}(2+a^2) \frac{1-b}{1+b} + \frac{1}{6}(1-a^2) \left(\frac{1-b}{1+b} \right)^2 \right] + \mathcal{O}(1) \end{aligned} \quad (5.8)$$

with $b^2 \equiv 3 \frac{2-a^2}{2+a^2}$ and a singular outcome for $\mathcal{O}(1)$. The one-loop correction yields

$$\frac{1}{2} \text{tr}' \ln S_N'' = \frac{1}{2} \eta \ln N - 2a \left[1 + \ln \frac{\pi}{2} - \ln \frac{2-a^2}{a} - 2(c^{-1} \arctan c - 1) \right] \quad (5.9)$$

where $c^2 \equiv 2 \frac{1-a^2}{2+a^2}$, which does not carry much meaning in view of the singular

* It is useful to remember that $\frac{1}{\pi} \int_{-1}^{+1} dx \frac{\sqrt{1-x^2}}{x^2 \pm a^2} = -1 + \Theta(a^2-1) \sqrt{1 \pm a^{-2}}$.

classical contribution at the same order.

The comparison of these semiclassical “results” with the exact formula (5.2) is poor. Although the leading terms agree (as is well-known) we have failed to reproduce the subleading contributions. The logarithmic N -dependence may still be restored by taking $\ln(x-y)^2 = \ln N^{-2}$ when $|x-y| < N^{-1}$, and by setting $\eta = -\frac{5}{6}$. The $\mathcal{O}(N)$ term is (incorrectly) a -dependent and even at $g=0$ ($a^2=1$) does not quite match up. We do, however, expect corrections of $\mathcal{O}(N)$ and below from a proper regularization of the one-loop determinant (see (4.9)), hence only the first line of eq. (5.8) can be trusted anyway. Presently, the first non-trivial subleading term of $F_N(g)-F_N(0)$ cannot be probed due to the divergencies, although it is interesting to note that the logarithmic singularity at $a^2=2$ already shows up correctly.

6. Conclusions

We have formulated the collective field theory for hermitean matrix models away from the large- N limit. By treating the Jacobian from the change of variables from eigenvalues x_i to their density $\rho(x)$ exactly, we arrived at a particular non-local and non-polynomial theory of two real scalar fields, ρ and λ . Nevertheless, we were able to set up a loop expansion for the free energy, at any $N < \infty$. The classical eigenvalue density, $\hat{\rho}$, is determined by a simple integral equation whose large- N limit becomes the usual airfoil equation. Its action naturally contains the entropy term for the Dyson gas of eigenvalues. The $\hbar$ expansion about the classical background yields sensible Feynman rules, including somewhat formal expressions for the one-loop determinant and the full propagator. It has to be stressed that the semiclassical (loop) expansion is *not* identical to the topological ($\frac{1}{N}$) expansion, because the “equation of motion” for $\hat{\rho}$ is not homogeneous in N , thanks to the Jacobian. We studied the collective field theory in some detail up to the three-loop level.

The attempt to develop our non-local field theory around $N = \infty$ met with three difficulties. First, the classical density $\hat{\rho}(x)$ can be expanded in $\frac{1}{N}$ only near the minimum of the matrix potential $V(x)$; elsewhere it behaves asymptotically as $e^{-NV(x)}$, a strong non-perturbative (in $\frac{1}{N}$) signal. The mixing of both regions due to the defining integral equation destroys the expansion everywhere; only the leading term $\hat{\rho} \rightarrow \rho_0$ is non-singular. Second, UV singularities $\propto (\text{tr } \mathbf{1})^k$ appear at k loops since the propagator is a delta-function to leading order. Amazingly, most of the divergent graphs cancel each other, except for the connected diagrams with the maximal number of loops and no self-contractions. As a result, the free energy is not just expanded in $\frac{1}{N}$ but also in $\frac{\delta(0)}{N\hat{\rho}}$. Third, the ensuing integrals $\int \hat{\rho}^{1-k}$ do not converge due to the exponentially small tail of $\hat{\rho}$, producing IR divergencies. A simple recipe to deal with these problems is to restrict the eigenvalue space to a finite interval and to implement a high-frequency cutoff. Preliminary answers were obtained here at the one-loop level.

Finally, a simple example was chosen to substantiate our abstract discussion and to compare with known results obtained by the method of orthogonal polynomials. As anticipated, only the planar contribution was reproduced entirely, but some features of the torus correction appeared as well.

Although we focused on the zero-dimensional hermitean one-matrix ensemble, the discussions in this paper should carry over *mutatis mutandis* to multi-matrix models, the one-dimensional theory, and to unitary matrix models.

It must be said that we did not yet achieve our goal of deriving a practical algorithm which systematically computes $\frac{1}{N}$ corrections in the semiclassical scheme. However, the necessary steps have been taken, and the details of our failure are no more puzzling as they are intriguing. We certainly hope to gain further insight into this outstanding problem in the future.

Acknowledgements: I wish to thank D. Karabali, J. Schiff and S. Wadia for useful discussions.

APPENDIX A

We give the basic notions of the infinite and finite Hilbert transforms, including some useful properties. Hilbert was the first to notice the (skew-)reciprocity of

$$u(x) = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{dy}{x-y} v(y) \quad , \quad v(x) = -\frac{1}{\pi} \int_{-\infty}^{\infty} \frac{dy}{x-y} u(y) \quad ; \quad (\text{A.1})$$

u is said to be conjugate to v . The (infinite) Hilbert transform

$$u(x) = \mathcal{H}_x[v] := \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{dy}{x-y} v(y) \quad (\text{A.2})$$

is related to Fourier integrals and to the Laplace transform and received an extensive treatment in Titchmarsh's book [13]. For $v \in L^{1+\epsilon}$, $\mathcal{H}[v]$ exists almost everywhere and also belongs to $L^{1+\epsilon}$, so its inverse is

$$v(x) \equiv \mathcal{H}_x^{-1}[u] = -\mathcal{H}_x[u] \quad . \quad (\text{A.3})$$

If $v \in L$, $\mathcal{H}[v]$ still exists almost everywhere but does not necessarily belong to L . We may, however, set $\mathcal{H}[1] = 0$. Furthermore, it is easily seen that

$$\mathcal{H}_x[v(ay+b)] = \frac{a}{|a|} \mathcal{H}_{ax+b}[v(y)] \quad . \quad (\text{A.4})$$

Some interesting properties of $\mathcal{H}$ generalize to the *finite* Hilbert transform,

$$u(x) = \mathcal{T}_x[v] := \frac{1}{\pi} \int_a^b \frac{dy}{x-y} v(y) \quad , \quad (\text{A.5})$$

which plays an important role in aerodynamics and has been investigated by a

number of authors [11, 12]. Most useful of those is Tricomi's convolution theorem,

$$\mathcal{T}\{v_1 \mathcal{T}[v_2] + v_2 \mathcal{T}[v_1]\} = \mathcal{T}[v_1] \mathcal{T}[v_2] - v_1 v_2 \quad , \quad (\text{A.6})$$

valid for $v_i \in L^{p_i}$ with $p_i > 1$ and $\frac{1}{p_1} + \frac{1}{p_2} < 1$. It is equivalent to the interchange formula

$$\int_a^b dy \int_a^b dz \frac{v_1(y) v_2(z)}{(x-y)(y-z)} = \int_a^b dz \int_a^b dy \frac{v_1(y) v_2(z)}{(x-y)(y-z)} - \pi^2 v_1(x) v_2(x) \quad (\text{A.7})$$

of Hardy and Poincaré (more generally, $v_1(y) v_2(z) \rightarrow F(x, y, z)$). Under the same conditions we also have Parseval's formula

$$\int_a^b dx \int_a^b dy \frac{v_1(x) v_2(y)}{x-y} = \int_a^b dy \int_a^b dx \frac{v_1(x) v_2(y)}{x-y} \quad (\text{A.8})$$

which may also be written as

$$\int_a^b dx \left(v_1 \mathcal{T}[v_2] + v_2 \mathcal{T}[v_1] \right) = 0 \quad , \quad (\text{A.9})$$

demonstrating the skew-adjointness of the $\mathcal{T}$ (and $\mathcal{H}$) operations with respect to the standard scalar product. Another convenient relation is

$$\mathcal{T}_x'[v] = \mathcal{T}_x[v'] \quad (\text{A.10})$$

which allows us to commute a derivative past the transform, as in (4.5). All these relations, (A.6)-(A.10), hold for $\mathcal{H}$ as well.

Conversely, the linearity property (A.4) cannot hold for $\mathcal{T}$; rather

$$\mathcal{T}_x[1] = -\frac{1}{\pi} \ln \left| \frac{x-b}{x-a} \right|, \quad (\text{A.11})$$

so that (A.6) with $v_1 = 1$ yields $\mathcal{T}\{\mathcal{T}[v]\} \neq -v$, in contradistinction to $\mathcal{H}$. This slightly complicates the inversion of $\mathcal{T}$. However, it is easily calculated that

$$\mathcal{T}_x \left[\frac{1}{\sqrt{(b-y)(y-a)}} \right] = 0, \quad \mathcal{T}_x \left[\sqrt{(b-y)(y-a)} \right] = x - \frac{a+b}{2} \quad (\text{A.12})$$

on the interval $x \in (a, b)$.[★] With these data a clever application of the convolution theorem (A.6) then establishes [12]

$$\mathcal{T}_x^{-1}[u] = \frac{1}{\sqrt{(b-x)(x-a)}} \left[\gamma - \frac{1}{\pi} \int_a^b \frac{dy}{x-y} u(y) \sqrt{(b-y)(y-a)} \right] \quad (\text{A.13})$$

where, using (A.8) and (A.12),

$$\gamma = \frac{1}{\pi} \int_a^b dx \mathcal{T}_x^{-1}[u] \quad (\text{A.14})$$

is an arbitrary normalization constant. The conditions of the convolution theorem restrict the validity of the inversion (A.13) to $u \in L^{\frac{4}{3}+\epsilon}$, but the uniqueness of the kernel (A.12) holds in $L^{1+\epsilon}$. For $u \in L^{1+\epsilon}$ the *r.h.s.* may not be in the same class; on the other side, any inverse of $\mathcal{T}$ in $L^{1+\epsilon}$ must be of the form (A.13). Furthermore, given $u \in L^{2+\epsilon}$, $\mathcal{T}[u]$ is necessarily orthogonal to the zero mode,

$$\int_a^b dx \frac{1}{\sqrt{(b-x)(x-a)}} \mathcal{T}_x[u] = 0 \quad (\text{A.15})$$

by virtue of the skew-adjointedness (A.9). This allows for alternative representa-

★ A note of caution: outside (a, b) the *r.h.s.* acquire an additional term of $\pm \epsilon [(x-b)(x-a)]^{\mp \frac{1}{2}}$, respectively, with $\epsilon = \Theta(x-b) - \Theta(a-x) = \pm 1$.

tions, *e.g.*

$$\mathcal{T}_x^{-1}[u] = \frac{\gamma'}{\sqrt{(b-x)(x-a)}} - \frac{1}{\pi} \int_a^b \frac{dy}{x-y} u(y) \frac{\sqrt{(b-x)(x-a)}}{\sqrt{(b-y)(y-a)}} \quad (\text{A.16})$$

which also goes to (A.3) when $(a, b) \rightarrow (-\infty, \infty)$.

Unfortunately, our nonlinear extension of the Hilbert transform,

$$u(x) = \mathcal{H}_x[v] - \beta \frac{v'}{v}(x) \quad (\text{A.17})$$

does not seem to have been investigated in the literature.

APPENDIX B

In the case of a gaussian ensemble ($V = \frac{1}{2}x^2$) of hermitean random matrices the free energy F_N for any N can be found *exactly*, with orthogonal polynomial techniques. As usual [4]

$$e^{-F_N} := \left[\prod_{i=1}^N \int dx_i \right] e^{-\frac{1}{2}N \sum_i x_i^2} \prod_{i < j} (x_i - x_j)^2 = N! \prod_{k=0}^{N-1} h_k \quad (\text{B.1})$$

where h_k is the norm of the k th monic orthogonal polynomial, $P_k(x) = x^k + \dots$. For our quadratic potential the P_k are just proportional to the Hermite polynomials,

$$P_k(x) = \sqrt{2N}^{-k} H_k(x\sqrt{N/2}) \quad , \quad (\text{B.2})$$

whose norms are well-known,

$$h_k = \int_{-\infty}^{+\infty} dx e^{-\frac{1}{2}Nx^2} P_k(x)^2 = N^{-k-\frac{1}{2}} k! \sqrt{2\pi} \quad . \quad (\text{B.3})$$

Hence, we find

$$e^{-F_N} = N^{-N^2/2} (2\pi)^{N/2} \prod_{k=1}^N k! \quad (\text{B.4})$$

which is the exact result.

We go on to expand

$$F_N = \frac{1}{2}N^2 \ln N - \frac{1}{2}N \ln 2\pi - \sum_1^N (N-k+1) \ln k \quad (\text{B.5})$$

in powers of $\frac{1}{N}$, with the help of Stirling's asymptotic formula[★]

$$\begin{aligned} \ln \Gamma(z) &= (z - \frac{1}{2}) \ln z - z + \frac{1}{2} \ln 2\pi + \sum_1^{s-1} \frac{B_{2k}}{2k(2k-1)} z^{1-2k} + R_s(z) \\ |R_s(z)| &< \frac{|B_{2s}|}{2s(2s-1) |z|^{2s-1} \cos^{2s-1}(\frac{1}{2} \arg z)} \end{aligned} \quad (\text{B.6})$$

for large values of $|z|$ as well as

$$\sum_{k=1}^N k \ln k = \int_0^{N+1} dt \ln \Gamma(t) + \frac{1}{2}(N+1)(N - \ln 2\pi) \quad (\text{B.7})$$

$$\int_p^{p+1} dt \ln \Gamma(t) = p \ln p - p + \frac{1}{2} \ln 2\pi \quad . \quad (\text{B.8})$$

The $N^2 \ln N$ terms cancel, and after collecting all the pieces we arrive at the asymptotic series

$$F_N \simeq \frac{3}{4}N^2 - N \ln N + (1 - \ln 2\pi)N - \frac{5}{12} \ln N + c_0 - \sum_{k=2}^{\infty} \frac{B_{2k} N^{2-2k}}{2k(2k-1)(2k-2)} \quad (\text{B.9})$$

where

$$\begin{aligned} c_0 &\simeq -\frac{1}{2} \ln 2\pi - \frac{1}{12} + \sum_2^{p-1} k \ln k \\ &\quad + \frac{1}{4}p^2 - \frac{1}{2}(p^2 - p + \frac{1}{6}) \ln p + \sum_2^{\infty} \frac{B_{2k} p^{2-2k}}{2k(2k-1)(2k-2)} \\ &= -0.7535173895042 \dots \end{aligned} \quad (\text{B.10})$$

is a universal constant, defined ever more accurately for ever larger values of the cut-off parameter p . As expected on general grounds, this is an expansion in $\frac{1}{N^2}$, except for a few terms at low orders.

★ B_{2k} denotes the Bernoulli numbers: $B_2 = \frac{1}{6}$, $B_4 = -\frac{1}{30}$, etc.

APPENDIX C

We outline a rough estimate of the determinant of the fluctuation operator (3.3), regulated by an IR cutoff L and a UV mode cutoff M . In effect, we restrict the eigenvalue range to $[-\frac{L}{2}, +\frac{L}{2}]$ with open boundary conditions and convert to Fourier modes

$$\rho(x) = \frac{1}{L} \sum_{k \in \mathbf{Z}} e^{2\pi i k x / L} \rho_k \quad (\text{C.1})$$

before setting $\rho_k = 0$ for $|k| > M$.

The Fourier transform of $S''_N(x, y)$ reads

$$S''_{kl} = \begin{pmatrix} -N^2 f_{kl} & -i\delta_{k+l,0} \\ -i\delta_{k+l,0} & \frac{1}{N} [\hat{\rho}_{k+l} - \hat{\rho}_k \hat{\rho}_l] \end{pmatrix} \quad (\text{C.2})$$

which, for the simple case of $V = \frac{1}{2}x^2$ and after reinstating $f(z) \rightarrow \ln z^2$, can be evaluated explicitly,

$$\begin{aligned} \hat{\rho}_k &= \frac{L}{2\pi k} J_1\left(\frac{4\pi k}{L}\right) \\ f_{kl} &= \frac{-2}{\pi k} \left(\text{si}(\pi k) + \frac{\pi}{2}\right) \delta_{k+l,0} \equiv f_k \delta_{k+l,0} \end{aligned} \quad (\text{C.3})$$

with $\text{si}(x) \equiv -\int_x^\infty \frac{\sin t}{t}$. By some algebraic identities we rewrite

$$\det' S'' = \det' [\delta_{k+l,0} - N f_k (\hat{\rho}_{k+l} - \hat{\rho}_k \hat{\rho}_l)] = \det [(\delta_{k+l,0} - \delta_{k,0} \delta_{l,0}) - N \sqrt{f_k} \hat{\rho}_{k+l} \sqrt{f_l}] \quad (\text{C.4})$$

where the prime means omission of the zero mode ($k=0$), and we have used $\hat{\rho}_0 = 1$.

For large values of $|k|$, one has

$$\hat{\rho}_k \approx \sqrt{\frac{L^3}{8\pi^4}} |k|^{-3/2} \cos\left(\frac{4\pi|k|}{L} - \frac{3\pi}{4}\right) \quad , \quad f_k \approx -\frac{1}{|k|} \quad (\text{C.5})$$

so that the matrix elements for (C.4) fall off as $|k+l|^{-5/2}$ perpendicular to the diagonal. We can make a rough estimate of $\det' S''$ by boldly taking into account

only the diagonal elements, *i.e.*

$$\det 'S'' \Big|_{\text{reg}} \approx -N f_0 \prod_{k=1}^M \left(1 + \frac{N}{k}\right)^2 = 2N \binom{M+N}{N}^2 \quad (\text{C.6})$$

since $f_0 = -2$. Let us now choose a UV cutoff proportional to N , say $M = \alpha N + \beta$.

Via Stirling's formula then

$$\frac{1}{2} \text{tr}' \ln S'' \Big|_{\text{reg}} \approx [(\alpha+1) \ln(\alpha+1) - \alpha \ln \alpha] N + \mathcal{O}(\ln N) \quad . \quad (\text{C.7})$$

More accurate estimates have to consider some sub-diagonals as well.

REFERENCES

1. M.L. Mehta: *Random matrices*, Academic Press (New York and London 1967).
2. E. Brézin and V.A. Kazakov, *Phys. Lett.* **B236** (1990), 144;
M.R. Douglas and S.H. Shenker, *Nucl. Phys.* **B335** (1990), 635;
D.J. Gross and A.A. Migdal, *Phys. Rev. Lett.* **64** (1990), 127,717;
E. Brézin, M.R. Douglas, V.A. Kazakov and S.H. Shenker, *Phys. Lett.* **B237** (1990), 43;
C. Crnković, P. Ginsparg and G. Moore, *Phys. Lett.* **B237** (1990), 196;
M.R. Douglas, *Phys. Lett.* **B238** (1990), 176.
3. E. Brézin, C. Itzykson, G. Parisi and J.B. Zuber, *Comm. Math. Phys.* **59** (1978), 35.
4. D. Bessis, C. Itzykson and J.B. Zuber, *Adv. Appl. Math.* **1** (1980), 109.
5. B. Sakita, *Quantum Theory of Many-Variable Systems and Fields*, World Scientific (Singapore 1985).
6. A. Jevicki, *Nucl. Phys.* **B146** (1978), 77.
7. A. Jevicki and B. Sakita, *Nucl. Phys.* **B185** (1981), 89.
8. J.D. Cohn and S.P. DeAlwis, IAS preprint IASSNS-HEP-91/7 (Feb. 1991).
9. D. Karabali and S. Sakita, *Int. J. Mod. Phys* **A6** (1991), 5079.
10. S. Ben-Menahem, SLAC preprint SLAC-PUB-5377 (Nov. 1990).
11. H. Söhlngen, *Math. Z.* **45** (1939), 245.
12. F.G. Tricomi, *Quart. J. Math. Oxford Ser. (2)* **2** (1951), 199.
13. E.C. Titchmarsh, *Introduction to the theory of Fourier integrals* (Oxford, 2nd ed. 1948) Chap. V.
14. O. Lechtenfeld, IAS preprint IASSNS-HEP-91/2 (Jan. 1990), to appear in *Journ. Mod. Phys.* **A**.